

# BiteMD™

External Analgesic  
First Aid Antiseptic



from the  
makers of

**Cutter**

## Insect Bite Relief



***Stops Pain***



***Relieves Itching***



***Reduces Infection***



Chiggers • Wasps • Mosquitoes • Ticks



***Soft Tip For  
Controlled Application***

Net 0.5 fl oz (14.8 mL)

from the  
makers of  
**Cutter**



External Analgesic  
First Aid Antiseptic

**BiteMD**




# BiteMD™

**External Analgesic  
First Aid Antiseptic**



from the makers of  
**Cutter**



*BiteMD, when applied soon after an insect bite, begins to work immediately to counteract the effects of bites and stings from bees, wasps, mosquitoes, ticks, hornets, ants, fire ants, deerflies, chiggers, sand fleas and horseflies.*

## DRUG FACTS:

### Active Ingredients

Benzocaine 5%  
Phenol 1.35%

**Purpose**  
Analgesic, Anesthetic, Antipruritic  
First Aid Antiseptic

**Uses:** First aid for temporary relief of:

- Pain
- Itching
- Infection

**Warnings:** For external use only

### When using this product

- Do not get in eyes.
- Do not apply over large areas of the body or bandage.
- In case of deep or puncture wounds, animal bites or serious burns, consult a doctor.

**Stop use and ask doctor** if condition worsens; or if symptoms persist for more than 7 days, clear up and occur again within a few days. Do not use longer than one week unless directed by doctor.

**Keep out of reach of children.** If swallowed, get medical help or contact a Poison Control Center right away. If in eyes, flush with water for 15 minutes and call a physician.

### Directions

- Clean area.
- Apply to the bite area by pressing straight down repeatedly until liquid flows, then dab sparingly.
- If itching persists, apply again in 10 to 15 minutes.
- For ticks and bees, remove tick or stinger before treatment.

**Adult and children  
2 years and over**

Apply to affected area  
1 to 3 times daily

**Children under 2 years**

Consult a doctor

**Inactive Ingredients** Camphor, Propylene Glycol

**IMPORTANT:** Read all information on this packaging and keep for future reference. This contains information not included on the stick label.

Manufactured by

Wisconsin Pharmacal Company, LLC

Jackson, WI 53037 USA

Distributed by Spectrum

Division of United Industries Corporation

PO Box 142642, St. Louis, MO 63114-0642

**Questions or Comments?** Call 1-800-767-9927.

HG-18-5253 ©2006 UIC

FPO UPC  
0 71121 95614 9

*For complete warnings and directions, see blister card packaging.*

Manufactured by Wisconsin Pharmacal Company, LLC, Jackson, WI 53037 USA

Distributed by Spectrum, Division of United Industries Corporation

PO Box 142642, St. Louis, MO 63114-0642

**Questions or Comments?** Call 1-800-767-9927.      HG-17-11777      ©2006 UIC

**BiteMD™**

External Analgesic  
First Aid Antiseptic



from the  
makers of **Cutter®**

Net 0.5 fl oz (14.8 mL)

Active Ingredients: Benzocaine 5%, Phenol 1.35%

**DIRECTIONS:** Adults and children 2 years and over – apply to bite area by pressing stick straight down repeatedly until liquid flows, then dab sparingly. If itching persists, apply again in 15 minutes. Apply to affected area 1 to 3 times daily. For ticks and bees, remove tick or stinger before treatment.

**WARNING:** For external use only. • Do not get in eyes.